

Oxygen (O₂) - Public Comment August 2023

ACTION: Medical Gas

- Essential for cellular metabolism and survival.
- First drug used for respiratory compromise or any time hypoxia is possible.

INDICATIONS:

- All patients with cardiopulmonary emergencies.
- Respiratory emergencies, including any patient who complains of shortness of breath.
- Chest pain.
- Suspected hypoxemia.

CONTRAINDICATIONS:

- Do NOT withhold oxygen from anyone who might need it.

ADULT DOSE/ROUTE:

Nasal Cannula: 2 to 6 liters/min (delivers 24 to 44% oxygen). **15 LPM for hyperoxygenation during endotracheal intubation**

Nonrebreather Mask: 10 to 15 liters/min (delivers 85 to 95% oxygen).

⇒ **BVM with O₂ Reservoir:** 15 to 25 liters/min (delivers 85 to 95% oxygen).

PEDIATRIC DOSE/ROUTE:

Nasal Cannula: 2 to 4 liters/min (delivers 24 to 38% oxygen). **15 LPM for hyperoxygenation during endotracheal intubation**

Nonrebreather Mask: 10 to 12 liters/min (delivers 85 to 95% oxygen).

BVM with O₂ Reservoir: 10 to 15 liters/min (delivers 60 to 95% oxygen).

NOTES:

- Target oxygen saturation levels when administering O₂ is 94 – 95%.
- Never withhold oxygen from anyone who might need it.
- Observe COPD patients closely and be prepared to ventilate them with BVM if necessary (development of apnea or increasing signs of respiratory failure).

Effective: xxxxxx
Supersedes: 03/01/15

DRAFT