


8.10 PEDIATRIC SEIZURE PUBLIC COMMENT JULY 2024

BLS – FAQ Link

- Assess **Vital Signs**,
- ABC's and responsiveness,
- **NPO**,
- **Oxygen** as needed
- Transport in left lateral recumbent position if no C-spine injury suspected

**DRAFT
VERSION**



Make Base Hospital Contact
Additional **Midazolam** administration beyond max dose needed for patient with continued seizures.

8.10 PEDIATRIC SEIZURE

PUBLIC COMMENT JULY 2024

BLS Treatment
<ul style="list-style-type: none"> • Assess airway, breathing, circulation and responsiveness. • Assess signs • Oxygen as indicated. • Transport in left lateral recumbent position if no C-spine injury is suspected.
ALS Treatment
<ul style="list-style-type: none"> • IV or IO of Normal Saline at TKO. • Check blood glucose. If blood glucose <60 mg/dl for children or <45mg/dl for neonates: administer Dextrose. <ul style="list-style-type: none"> ○ Neonates 1 month: D10W, 5ml/kg IV/IO (0.5g/kg. Max 25 gram) ○ Children >1 month: D10W, 5ml/kg IV/IO (0.5g/kg. Max 25 gram) • If no IV or IO access: administer Glucagon. <ul style="list-style-type: none"> ⊖ Less than 20kg: 0.5mg IM/IV ○ Greater than 20kg: 1mg IM/IV • If status epilepticus is present, administer Midazolam or Diastat rectal gel: • Status epilepticus is continuous seizure activity lasting > 5 minutes OR multiple seizures without regaining consciousness between seizures. • Status epilepticus is continuous seizure activity lasting > 5 minutes OR multiple seizures without regaining consciousness between seizures. <ul style="list-style-type: none"> • Midazolam: <ul style="list-style-type: none"> • 0.2 mg/kg IM x1 (Max 10mg) • 0.2 mg/kg intranasally x1 (half in each nostril) (Max 10mg) • 0.1 mg/kg slow push IV/IO (max 5mg). May repeat IV/IO dose in 5 min x1. Maximum total dose 0.2mg/kg IV/IO (not to exceed 10mg) • Administer Midazolam IV only if IV in place when seizure occurs. • Diastat rectal gel: Preferred in patients who are on the medication, it is readily available, and the dosage is decipherable on the container. May assist caregiver in administering normally prescribed at home medication
Base Hospital Contact
Additional Midazolam administration beyond max dose needed for patient with continued seizures.

DRAFT