

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through December 2019

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (628) 217-6336, FAX (415) 431-0353

Director of Health: Grant Colfax, MD
Director of Population Health Division: Tomas Aragon, MD, DrPH
Director of ARCHES: Wayne Enanoria, PhD, MPH
Directors of HIV Surveillance Section: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 12/31/2019)

Persons living with HIV infection ²	15,908
Diagnoses of HIV infection stage 3 (AIDS), cumulative	30,167
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	21,114

California³ (as of 12/31/2017)

Persons living with HIV infection ²	135,082
Diagnoses of HIV infection stage 3 (AIDS), cumulative	NA
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	NA

United States⁴ (as of 12/31/2017)

Persons living with HIV infection ²	1,020,419
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2018)	1,290,902
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	740,377


¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.

³ CA data source: California HIV Surveillance Report 2018
https://www.cdph.ca.gov/Programs/CID/DOA/Pages/OA_case_surveillance_reports.aspx


⁴ US data source: CDC HIV Surveillance Report 2018, vol. 30 (Table 2b, 19b, 22b).
<https://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2019*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2019 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2008-2019

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	4 (0.8)	1 (0.2)	3 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	56 (10.8)	65 (13.7)	59 (12.6)	45 (10.5)	65 (13.9)	59 (14.8)
25 - 29	81 (15.6)	66 (13.9)	66 (14.0)	69 (16.1)	75 (16.1)	79 (19.8)
30 - 39	179 (34.4)	141 (29.6)	146 (31.1)	117 (27.3)	141 (30.2)	115 (28.8)
40 - 49	152 (29.2)	122 (25.6)	129 (27.4)	131 (30.5)	131 (28.1)	99 (24.8)
50 - 59	38 (7.3)	64 (13.4)	51 (10.9)	52 (12.1)	43 (9.2)	35 (8.8)
60 +	10 (1.9)	17 (3.6)	16 (3.4)	15 (3.5)	12 (2.6)	13 (3.3)
Total	520 (100)	476 (100)	470 (100)	429 (100)	467 (100)	400 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.3)	3 (1.0)	0 (0.0)	3 (1.3)	1 (0.5)	0 (0.0)
18 - 24	39 (12.1)	37 (12.8)	34 (14.4)	24 (10.3)	30 (14.9)	12 (7.4)
25 - 29	58 (18.0)	71 (24.7)	53 (22.5)	41 (17.6)	38 (18.8)	26 (16.0)
30 - 39	96 (29.7)	87 (30.2)	76 (32.2)	79 (33.9)	65 (32.2)	63 (38.7)
40 - 49	75 (23.2)	59 (20.5)	37 (15.7)	48 (20.6)	42 (20.8)	28 (17.2)
50 - 59	41 (12.7)	18 (6.3)	24 (10.2)	27 (11.6)	16 (7.9)	19 (11.7)
60 +	13 (4.0)	13 (4.5)	12 (5.1)	11 (4.7)	10 (5.0)	15 (9.2)
Total	323 (100)	288 (100)	236 (100)	233 (100)	202 (100)	163 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2008-2019

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	17 (3.9)	15 (4.6)	15 (5.0)	12 (4.8)	11 (4.5)	10 (5.3)
25 - 29	31 (7.1)	23 (7.1)	26 (8.7)	21 (8.3)	25 (10.3)	17 (8.9)
30 - 39	123 (28.3)	75 (23.1)	74 (24.7)	62 (24.6)	60 (24.7)	43 (22.6)
40 - 49	167 (38.4)	120 (37.0)	99 (33.1)	99 (39.3)	83 (34.2)	60 (31.6)
50 - 59	75 (17.2)	69 (21.3)	62 (20.7)	49 (19.4)	44 (18.1)	39 (20.5)
60 +	21 (4.8)	21 (6.5)	23 (7.7)	9 (3.6)	20 (8.2)	21 (11.1)
Total	435 (100)	324 (100)	299 (100)	252 (100)	243 (100)	190 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.8)	0 (0.0)	0 (0.0)
18 - 24	5 (3.6)	4 (3.2)	2 (2.1)	1 (0.8)	3 (2.6)	2 (1.9)
25 - 29	13 (9.4)	9 (7.1)	8 (8.2)	15 (11.6)	12 (10.5)	8 (7.6)
30 - 39	41 (29.5)	35 (27.8)	27 (27.8)	31 (24.0)	32 (28.1)	31 (29.5)
40 - 49	40 (28.8)	36 (28.6)	23 (23.7)	41 (31.8)	21 (18.4)	24 (22.9)
50 - 59	27 (19.4)	28 (22.2)	19 (19.6)	14 (10.9)	26 (22.8)	22 (21.0)
60 +	13 (9.4)	14 (11.1)	18 (18.6)	26 (20.2)	20 (17.5)	18 (17.1)
Total	139 (100)	126 (100)	97 (100)	129 (100)	114 (100)	105 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2008-2019

Gender (2)	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Men	461 (88.7)	423 (88.9)	417 (88.7)	375 (87.4)	432 (92.5)	366 (91.5)
Women	41 (7.9)	31 (6.5)	36 (7.7)	41 (9.6)	23 (4.9)	22 (5.5)
Trans Women	18 (3.5)	22 (4.6)	17 (3.6)	13 (3.0)	12 (2.6)	12 (3.0)
Total	520 (100)	476 (100)	470 (100)	429 (100)	467 (100)	400 (100)

Gender (2)	Year of initial HIV diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
Men	298 (92.3)	253 (87.8)	202 (85.6)	200 (85.8)	177 (87.6)	137 (84.0)
Women	13 (4.0)	28 (9.7)	25 (10.6)	27 (11.6)	17 (8.4)	13 (8.0)
Trans Women	12 (3.7)	7 (2.4)	8 (3.4)	6 (2.6)	7 (3.5)	12 (7.4)
Total	323 (100)	288 (100)	236 (100)	233 (100)	202 (100)	163 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Data on trans men are not released separately due to small numbers.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2008-2019

Gender (2)	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Men	375 (86.2)	288 (88.9)	267 (89.3)	214 (84.9)	210 (86.4)	167 (87.9)
Women	43 (9.9)	24 (7.4)	16 (5.4)	24 (9.5)	24 (9.9)	17 (8.9)
Trans Women	16 (3.7)	12 (3.7)	16 (5.4)	14 (5.6)	9 (3.7)	6 (3.2)
Total	435 (100)	324 (100)	299 (100)	252 (100)	243 (100)	190 (100)

Gender (2)	Year of AIDS diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
Men	125 (89.9)	99 (78.6)	83 (85.6)	101 (78.3)	99 (86.8)	93 (88.6)
Women	7 (5.0)	19 (15.1)	11 (11.3)	22 (17.1)	12 (10.5)	5 (4.8)
Trans Women	7 (5.0)	8 (6.3)	2 (2.1)	6 (4.7)	3 (2.6)	7 (6.7)
Total	139 (100)	126 (100)	97 (100)	129 (100)	114 (100)	105 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Data on trans men are not released separately due to small numbers.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2008-2019

Race/ethnicity	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
White	251 (48.3)	227 (47.7)	224 (47.7)	221 (51.5)	229 (49.0)	181 (45.3)
African American	77 (14.8)	63 (13.2)	64 (13.6)	65 (15.2)	50 (10.7)	48 (12.0)
Latinx	123 (23.7)	117 (24.6)	109 (23.2)	87 (20.3)	114 (24.4)	105 (26.3)
Asian/Pacific Islander/ Native American	45 (8.7)	43 (9.0)	42 (8.9)	38 (8.9)	57 (12.2)	51 (12.8)
Multiple races	24 (4.6)	25 (5.3)	30 (6.4)	17 (4.0)	16 (3.4)	14 (3.5)
Total (2)	520 (100)	476 (100)	470 (100)	429 (100)	467 (100)	400 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
White	136 (42.1)	119 (41.3)	90 (38.1)	85 (36.5)	59 (29.2)	58 (35.6)
African American	33 (10.2)	42 (14.6)	34 (14.4)	39 (16.7)	41 (20.3)	28 (17.2)
Latinx	95 (29.4)	75 (26.0)	69 (29.2)	66 (28.3)	78 (38.6)	51 (31.3)
Asian/Pacific Islander/ Native American	45 (13.9)	36 (12.5)	37 (15.7)	32 (13.7)	19 (9.4)	22 (13.5)
Multiple races	14 (4.3)	15 (5.2)	6 (2.5)	11 (4.7)	5 (2.5)	2 (1.2)
Total (2)	323 (100)	288 (100)	236 (100)	233 (100)	202 (100)	163 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2008-2019

Race/ethnicity	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
White	218 (50.1)	170 (52.5)	145 (48.5)	128 (50.8)	121 (49.8)	93 (48.9)
African American	79 (18.2)	60 (18.5)	49 (16.4)	46 (18.3)	40 (16.5)	31 (16.3)
Latinx	78 (17.9)	66 (20.4)	64 (21.4)	48 (19.0)	54 (22.2)	36 (18.9)
Asian/Pacific Islander/ Native American	39 (9.0)	15 (4.6)	23 (7.7)	14 (5.6)	15 (6.2)	18 (9.5)
Multiple races	21 (4.8)	13 (4.0)	18 (6.0)	16 (6.3)	13 (5.3)	12 (6.3)
Total (2)	435 (100)	324 (100)	299 (100)	252 (100)	243 (100)	190 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
White	52 (37.4)	61 (48.4)	45 (46.4)	52 (40.3)	36 (31.6)	46 (43.8)
African American	28 (20.1)	19 (15.1)	16 (16.5)	25 (19.4)	18 (15.8)	18 (17.1)
Latinx	32 (23.0)	32 (25.4)	20 (20.6)	35 (27.1)	41 (36.0)	19 (18.1)
Asian/Pacific Islander/ Native American	17 (12.2)	5 (4.0)	11 (11.3)	13 (10.1)	10 (8.8)	15 (14.3)
Multiple races	10 (7.2)	9 (7.1)	5 (5.2)	4 (3.1)	9 (7.9)	7 (6.7)
Total (2)	139 (100)	126 (100)	97 (100)	129 (100)	114 (100)	105 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2008-2019

Transmission category (2)	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Gay or bisexual male	368 (70.8)	330 (69.3)	307 (65.3)	304 (70.9)	360 (77.1)	309 (77.3)
Injection drug user (IDU)	34 (6.5)	26 (5.5)	39 (8.3)	28 (6.5)	21 (4.5)	21 (5.3)
Gay or bisexual male IDU	72 (13.8)	81 (17.0)	68 (14.5)	60 (14.0)	52 (11.1)	48 (12.0)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	36 (6.9)	26 (5.5)	38 (8.1)	30 (7.0)	26 (5.6)	17 (4.3)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	10 (1.9)	13 (2.7)	18 (3.8)	7 (1.6)	8 (1.7)	5 (1.3)
Total	520 (100)	476 (100)	470 (100)	429 (100)	467 (100)	400 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
Gay or bisexual male	242 (74.9)	218 (75.7)	165 (69.9)	143 (61.4)	130 (64.4)	109 (66.9)
Injection drug user (IDU)	25 (7.7)	17 (5.9)	21 (8.9)	26 (11.2)	26 (12.9)	13 (8.0)
Gay or bisexual male IDU	43 (13.3)	25 (8.7)	26 (11.0)	31 (13.3)	25 (12.4)	19 (11.7)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	11 (3.4)	21 (7.3)	17 (7.2)	16 (6.9)	11 (5.4)	9 (5.5)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	2 (0.6)	7 (2.4)	7 (3.0)	17 (7.3)	10 (5.0)	13 (8.0)
Total	323 (100)	288 (100)	236 (100)	233 (100)	202 (100)	163 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2008-2019

Transmission category (2)	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Gay or bisexual male	261 (60.0)	208 (64.2)	170 (56.9)	149 (59.1)	151 (62.1)	112 (58.9)
Injection drug user (IDU)	34 (7.8)	35 (10.8)	23 (7.7)	34 (13.5)	23 (9.5)	19 (10.0)
Gay or bisexual male IDU	99 (22.8)	56 (17.3)	83 (27.8)	48 (19.0)	43 (17.7)	46 (24.2)
Transfusion/Hemophilia	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	32 (7.4)	18 (5.6)	18 (6.0)	17 (6.7)	21 (8.6)	12 (6.3)
Perinatal	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.4)	0 (0.0)	0 (0.0)
Other/Unknown	9 (2.1)	5 (1.5)	5 (1.7)	3 (1.2)	5 (2.1)	1 (0.5)
Total	435 (100)	324 (100)	299 (100)	252 (100)	243 (100)	190 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
Gay or bisexual male	95 (68.3)	72 (57.1)	58 (59.8)	61 (47.3)	68 (59.6)	59 (56.2)
Injection drug user (IDU)	16 (11.5)	19 (15.1)	12 (12.4)	14 (10.9)	13 (11.4)	13 (12.4)
Gay or bisexual male IDU	21 (15.1)	23 (18.3)	16 (16.5)	29 (22.5)	21 (18.4)	26 (24.8)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	7 (5.0)	11 (8.7)	11 (11.3)	19 (14.7)	8 (7.0)	4 (3.8)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	0 (0.0)	1 (0.8)	0 (0.0)	6 (4.7)	4 (3.5)	3 (2.9)
Total	139 (100)	126 (100)	97 (100)	129 (100)	114 (100)	105 (100)

- (1) Data in recent years are incomplete due to delays in case reporting.
(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2019

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	656	348	13303
2003	651	332	13622
2004	691	337	13976
2005	566	346	14196
2006	540	326	14410
2007	536	324	14622
2008	520	263	14879
2009	476	251	15104
2010	470	244	15330
2011	429	236	15523
2012	467	236	15754
2013	400	259	15895
2014	323	239	15979
2015	288	256	16011
2016	236	236	16011
2017	233	254	15990
2018	202	256	15936
2019	163	191	15908

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2019

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	8	21
1982	99	32	88
1983	274	111	251
1984	557	273	535
1985	859	534	860
1986	1237	807	1290
1987	1636	878	2048
1988	1762	1038	2772
1989	2159	1279	3652
1990	2043	1364	4331
1991	2284	1512	5103
1992	2326	1639	5790
1993	2064	1603	6251
1994	1777	1600	6428
1995	1552	1485	6495
1996	1067	993	6569
1997	795	424	6940
1998	686	401	7225
1999	575	353	7447
2000	552	350	7649
2001	507	325	7831
2002	482	321	7992
2003	545	296	8241
2004	477	301	8417
2005	476	310	8583
2006	451	288	8746
2007	443	270	8919
2008	435	227	9127
2009	324	208	9243
2010	299	193	9349
2011	252	190	9411
2012	243	182	9472
2013	190	191	9471
2014	139	188	9422
2015	126	203	9345
2016	97	187	9255
2017	129	203	9181
2018	114	208	9087
2019	105	139	9053
Total	30167	21114	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	African American No. (%)	Latinx No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Multiple races No. (%)	Total (1) No. (%)
Men						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.2)	1 (0.0)
18 - 24	12 (0.1)	15 (1.0)	20 (0.7)	6 (0.6)	3 (0.7)	56 (0.4)
25 - 29	67 (0.8)	45 (3.1)	99 (3.5)	41 (4.3)	9 (2.0)	262 (1.8)
30 - 39	595 (6.7)	161 (11.2)	497 (17.4)	186 (19.5)	61 (13.5)	1500 (10.3)
40 - 49	1339 (15.1)	202 (14.1)	655 (23.0)	278 (29.1)	112 (24.8)	2589 (17.7)
50 - 59	3332 (37.5)	484 (33.7)	1013 (35.5)	286 (29.9)	158 (35.0)	5274 (36.2)
60 +	3540 (39.8)	529 (36.8)	568 (19.9)	158 (16.5)	107 (23.7)	4905 (33.6)
Men Total	8885 (100)	1436 (100)	2852 (100)	955 (100)	451 (100)	14587 (100)
Women						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	1 (0.3)	1 (0.5)	0 (0.0)	0 (0.0)	2 (0.2)
18 - 24	1 (0.4)	5 (1.5)	4 (1.9)	0 (0.0)	2 (6.1)	12 (1.3)
25 - 29	6 (2.4)	6 (1.8)	8 (3.9)	1 (1.4)	2 (6.1)	23 (2.6)
30 - 39	23 (9.1)	34 (10.2)	41 (19.8)	10 (13.7)	3 (9.1)	111 (12.3)
40 - 49	49 (19.4)	54 (16.2)	41 (19.8)	19 (26.0)	10 (30.3)	174 (19.3)
50 - 59	103 (40.9)	112 (33.6)	57 (27.5)	23 (31.5)	12 (36.4)	307 (34.1)
60 +	70 (27.8)	121 (36.3)	55 (26.6)	20 (27.4)	4 (12.1)	271 (30.1)
Women Total	252 (100)	333 (100)	207 (100)	73 (100)	33 (100)	900 (100)
Trans Women	74 (100)	128 (100)	148 (100)	43 (100)	21 (100)	414 (100)

(1) Total includes persons with unknown race. Data on trans men are not released separately due to small numbers.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2019

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White		African American		Latinx		Asian/ Pacific Islander/ Native American		Multiple races		Total (2)	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Men												
Gay or bisexual male	7266	(81.8)	874	(60.9)	2347	(82.3)	816	(85.4)	325	(72.1)	11630	(79.7)
Injection drug user (IDU)	170	(1.9)	180	(12.5)	72	(2.5)	21	(2.2)	17	(3.8)	460	(3.2)
Gay or bisexual male IDU	1362	(15.3)	272	(18.9)	335	(11.7)	79	(8.3)	99	(22.0)	2147	(14.7)
Transfusion/Hemophiliac	6	(0.1)	2	(0.1)	2	(0.1)	3	(0.3)	0	(0.3)	13	(0.0)
Heterosexual (3)	33	(0.4)	73	(5.1)	51	(1.8)	21	(2.2)	6	(1.3)	184	(1.3)
Perinatal	1	(0.0)	3	(0.2)	2	(0.1)	1	(0.1)	2	(0.4)	9	(0.1)
Other/Unknown	47	(0.5)	32	(2.2)	43	(1.5)	14	(1.5)	2	(0.4)	144	(1.0)
Men Total	8885	(100)	1436	(100)	2852	(100)	955	(100)	451	(100)	14587	(100)
Women												
Injection drug user (IDU)	146	(57.9)	162	(48.6)	62	(30.0)	13	(17.8)	22	(66.7)	405	(45.0)
Transfusion/Hemophilia	5	(2.0)	1	(0.3)	2	(1.0)	2	(2.7)	0	(0.0)	10	(1.1)
Heterosexual (3)	82	(32.5)	138	(41.4)	113	(54.6)	52	(71.2)	6	(18.2)	391	(43.4)
Perinatal	1	(0.4)	6	(1.8)	9	(4.3)	0	(0.0)	2	(6.1)	18	(2.0)
Other/Unknown	18	(7.1)	26	(7.8)	21	(10.1)	6	(8.2)	3	(9.1)	76	(8.4)
Women Total	252	(100)	333	(100)	207	(100)	73	(100)	33	(100)	900	(100)
Trans Women	74	(100)	128	(100)	148	(100)	43	(100)	21	(100)	414	(100)

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Total includes persons with unknown race. Data on trans men are not released separately due to small numbers.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.