

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through June 2019

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 437-6336, FAX (415) 431-0353

Director of Health: Grant Colfax, MD
Director of Population Health Division: Tomas Aragon, MD, DrPH
Director of ARCHES: Wayne Enanoria, PhD, MPH
Directors of HIV Surveillance Section: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 06/30/2019)

Persons living with HIV infection ²	15,957
Diagnoses of HIV infection stage 3 (AIDS), cumulative	30,131
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	21,013

California³ (as of 12/31/2017)

Persons living with HIV infection ²	135,082
Diagnoses of HIV infection stage 3 (AIDS), cumulative	NA
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	NA

United States⁴ (as of 12/31/2016)

Persons living with HIV infection ²	1,008,929
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2017)	1,281,787
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	729,796


¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.

³ CA data source: California HIV Surveillance Report 2017
https://www.cdph.ca.gov/Programs/CID/DOA/Pages/OA_case_surveillance_reports.aspx


⁴ US data source: CDC HIV Surveillance Report 2017, vol. 29 (Table 2b, 19b, 22b).
<https://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2019*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2019 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2008-2019

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	4 (0.8)	1 (0.2)	3 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	56 (10.8)	62 (13.1)	59 (12.6)	45 (10.6)	64 (13.7)	58 (14.5)
25 - 29	80 (15.4)	66 (14.0)	66 (14.0)	69 (16.2)	75 (16.1)	80 (20.1)
30 - 39	180 (34.6)	140 (29.7)	146 (31.1)	115 (27.0)	141 (30.3)	115 (28.8)
40 - 49	152 (29.2)	122 (25.8)	129 (27.4)	130 (30.5)	131 (28.1)	98 (24.6)
50 - 59	38 (7.3)	64 (13.6)	51 (10.9)	52 (12.2)	43 (9.2)	35 (8.8)
60 +	10 (1.9)	17 (3.6)	16 (3.4)	15 (3.5)	12 (2.6)	13 (3.3)
Total	520 (100)	472 (100)	470 (100)	426 (100)	466 (100)	399 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.3)	3 (1.1)	0 (0.0)	3 (1.3)	1 (0.5)	0 (0.0)
18 - 24	39 (12.1)	37 (13.2)	33 (14.0)	23 (10.1)	29 (14.6)	6 (7.4)
25 - 29	58 (18.0)	67 (23.8)	55 (23.4)	41 (18.0)	39 (19.6)	11 (13.6)
30 - 39	95 (29.5)	85 (30.2)	75 (31.9)	78 (34.2)	62 (31.2)	32 (39.5)
40 - 49	75 (23.3)	58 (20.6)	37 (15.7)	47 (20.6)	41 (20.6)	14 (17.3)
50 - 59	41 (12.7)	18 (6.4)	24 (10.2)	25 (11.0)	17 (8.5)	12 (14.8)
60 +	13 (4.0)	13 (4.6)	11 (4.7)	11 (4.8)	10 (5.0)	6 (7.4)
Total	322 (100)	281 (100)	235 (100)	228 (100)	199 (100)	81 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2008-2019

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	17 (3.9)	15 (4.6)	15 (5.0)	12 (4.8)	11 (4.5)	10 (5.3)
25 - 29	31 (7.1)	23 (7.1)	26 (8.7)	21 (8.3)	25 (10.3)	17 (8.9)
30 - 39	123 (28.3)	75 (23.1)	74 (24.7)	62 (24.6)	60 (24.7)	43 (22.6)
40 - 49	167 (38.4)	120 (37.0)	99 (33.1)	99 (39.3)	83 (34.2)	60 (31.6)
50 - 59	75 (17.2)	69 (21.3)	62 (20.7)	49 (19.4)	44 (18.1)	39 (20.5)
60 +	21 (4.8)	21 (6.5)	23 (7.7)	9 (3.6)	20 (8.2)	21 (11.1)
Total	435 (100)	324 (100)	299 (100)	252 (100)	243 (100)	190 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.8)	0 (0.0)	0 (0.0)
18 - 24	5 (3.6)	4 (3.2)	2 (2.1)	1 (0.8)	3 (2.6)	0 (0.0)
25 - 29	13 (9.4)	9 (7.1)	8 (8.2)	15 (11.6)	12 (10.5)	3 (6.4)
30 - 39	41 (29.5)	35 (27.8)	28 (28.9)	31 (24.0)	32 (28.1)	13 (27.7)
40 - 49	40 (28.8)	36 (28.6)	22 (22.7)	41 (31.8)	21 (18.4)	10 (21.3)
50 - 59	27 (19.4)	28 (22.2)	18 (18.6)	14 (10.9)	26 (22.8)	10 (21.3)
60 +	13 (9.4)	14 (11.1)	19 (19.6)	26 (20.2)	20 (17.5)	11 (23.4)
Total	139 (100)	126 (100)	97 (100)	129 (100)	114 (100)	47 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2008-2019

Gender (2)	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Men	461 (88.7)	420 (89.0)	417 (88.7)	372 (87.3)	431 (92.5)	365 (91.5)
Women	41 (7.9)	30 (6.4)	36 (7.7)	41 (9.6)	23 (4.9)	22 (5.5)
Trans Women	18 (3.5)	22 (4.7)	17 (3.6)	13 (3.1)	12 (2.6)	12 (3.0)
Total	520 (100)	472 (100)	470 (100)	426 (100)	466 (100)	399 (100)

Gender (2)	Year of initial HIV diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
Men	297 (92.2)	247 (87.9)	202 (86.0)	195 (85.5)	174 (87.4)	68 (84.0)
Women	13 (4.0)	27 (9.6)	24 (10.2)	27 (11.8)	17 (8.5)	7 (8.6)
Trans Women	12 (3.7)	7 (2.5)	8 (3.4)	6 (2.6)	7 (3.5)	6 (7.4)
Total	322 (100)	281 (100)	235 (100)	228 (100)	199 (100)	81 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Data on trans men are not released separately due to small numbers.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2008-2019

Gender (2)	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Men	375 (86.2)	288 (88.9)	267 (89.3)	214 (84.9)	210 (86.4)	167 (87.9)
Women	43 (9.9)	24 (7.4)	16 (5.4)	24 (9.5)	24 (9.9)	17 (8.9)
Trans Women	16 (3.7)	12 (3.7)	16 (5.4)	14 (5.6)	9 (3.7)	6 (3.2)
Total	435 (100)	324 (100)	299 (100)	252 (100)	243 (100)	190 (100)

Gender (2)	Year of AIDS diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
Men	125 (89.9)	99 (78.6)	84 (86.6)	101 (78.3)	99 (86.8)	43 (91.5)
Women	7 (5.0)	19 (15.1)	10 (10.3)	22 (17.1)	12 (10.5)	2 (4.3)
Trans Women	7 (5.0)	8 (6.3)	2 (2.1)	6 (4.7)	3 (2.6)	2 (4.3)
Total	139 (100)	126 (100)	97 (100)	129 (100)	114 (100)	47 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Data on trans men are not released separately due to small numbers.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2008-2019

Race/ethnicity	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
White	252 (48.5)	226 (47.9)	225 (47.9)	220 (51.6)	229 (49.1)	181 (45.4)
African American	78 (15.0)	63 (13.3)	63 (13.4)	65 (15.3)	50 (10.7)	48 (12.0)
Latino	121 (23.3)	115 (24.4)	109 (23.2)	85 (20.0)	113 (24.2)	104 (26.1)
Asian/Pacific Islander/ Native American	45 (8.7)	41 (8.7)	42 (8.9)	38 (8.9)	57 (12.2)	51 (12.8)
Multiple races	24 (4.6)	26 (5.5)	30 (6.4)	17 (4.0)	16 (3.4)	14 (3.5)
Total (2)	520 (100)	472 (100)	470 (100)	426 (100)	466 (100)	399 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
White	135 (41.9)	114 (40.6)	90 (38.3)	83 (36.4)	58 (29.1)	33 (40.7)
African American	33 (10.2)	41 (14.6)	34 (14.5)	38 (16.7)	41 (20.6)	9 (11.1)
Latino	95 (29.5)	74 (26.3)	70 (29.8)	64 (28.1)	75 (37.7)	25 (30.9)
Asian/Pacific Islander/ Native American	45 (14.0)	36 (12.8)	35 (14.9)	32 (14.0)	20 (10.1)	12 (14.8)
Multiple races	14 (4.3)	15 (5.3)	6 (2.6)	11 (4.8)	5 (2.5)	1 (1.2)
Total (2)	322 (100)	281 (100)	235 (100)	228 (100)	199 (100)	81 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2008-2019

Race/ethnicity	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
White	218 (50.1)	170 (52.5)	144 (48.2)	128 (50.8)	121 (49.8)	93 (48.9)
African American	79 (18.2)	60 (18.5)	49 (16.4)	46 (18.3)	40 (16.5)	32 (16.8)
Latino	78 (17.9)	66 (20.4)	64 (21.4)	48 (19.0)	54 (22.2)	35 (18.4)
Asian/Pacific Islander/ Native American	39 (9.0)	15 (4.6)	23 (7.7)	14 (5.6)	15 (6.2)	18 (9.5)
Multiple races	21 (4.8)	13 (4.0)	19 (6.4)	16 (6.3)	13 (5.3)	12 (6.3)
Total (2)	435 (100)	324 (100)	299 (100)	252 (100)	243 (100)	190 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
White	52 (37.4)	61 (48.4)	46 (47.4)	52 (40.3)	35 (30.7)	20 (42.6)
African American	28 (20.1)	19 (15.1)	15 (15.5)	25 (19.4)	18 (15.8)	8 (17.0)
Latino	32 (23.0)	32 (25.4)	20 (20.6)	35 (27.1)	41 (36.0)	6 (12.8)
Asian/Pacific Islander/ Native American	17 (12.2)	5 (4.0)	11 (11.3)	13 (10.1)	10 (8.8)	6 (12.8)
Multiple races	10 (7.2)	9 (7.1)	5 (5.2)	4 (3.1)	10 (8.8)	7 (14.9)
Total (2)	139 (100)	126 (100)	97 (100)	129 (100)	114 (100)	47 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report

Cases Reported as of 06/30/2019

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2008-2019

Transmission category (2)	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Gay or bisexual male	367 (70.6)	327 (69.3)	307 (65.3)	302 (70.9)	359 (77.0)	308 (77.2)
Injection drug user (IDU)	34 (6.5)	26 (5.5)	39 (8.3)	28 (6.6)	21 (4.5)	21 (5.3)
Gay or bisexual male IDU	73 (14.0)	81 (17.2)	68 (14.5)	59 (13.8)	52 (11.2)	48 (12.0)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	36 (6.9)	25 (5.3)	38 (8.1)	30 (7.0)	26 (5.6)	17 (4.3)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	10 (1.9)	13 (2.8)	18 (3.8)	7 (1.6)	8 (1.7)	5 (1.3)
Total	520 (100)	472 (100)	470 (100)	426 (100)	466 (100)	399 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
Gay or bisexual male	242 (75.2)	212 (75.4)	166 (70.6)	139 (61.0)	127 (63.8)	53 (65.4)
Injection drug user (IDU)	25 (7.8)	17 (6.0)	21 (8.9)	26 (11.4)	26 (13.1)	9 (11.1)
Gay or bisexual male IDU	42 (13.0)	25 (8.9)	25 (10.6)	31 (13.6)	25 (12.6)	9 (11.1)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	11 (3.4)	20 (7.1)	16 (6.8)	16 (7.0)	11 (5.5)	4 (4.9)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	2 (0.6)	7 (2.5)	7 (3.0)	16 (7.0)	10 (5.0)	6 (7.4)
Total	322 (100)	281 (100)	235 (100)	228 (100)	199 (100)	81 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2008-2019

Transmission category (2)	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Gay or bisexual male	261 (60.0)	208 (64.2)	170 (56.9)	149 (59.1)	151 (62.1)	112 (58.9)
Injection drug user (IDU)	34 (7.8)	35 (10.8)	23 (7.7)	34 (13.5)	23 (9.5)	19 (10.0)
Gay or bisexual male IDU	99 (22.8)	56 (17.3)	83 (27.8)	48 (19.0)	43 (17.7)	46 (24.2)
Transfusion/Hemophilia	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	32 (7.4)	18 (5.6)	18 (6.0)	17 (6.7)	21 (8.6)	12 (6.3)
Perinatal	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.4)	0 (0.0)	0 (0.0)
Other/Unknown	9 (2.1)	5 (1.5)	5 (1.7)	3 (1.2)	5 (2.1)	1 (0.5)
Total	435 (100)	324 (100)	299 (100)	252 (100)	243 (100)	190 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)
Gay or bisexual male	96 (69.1)	72 (57.1)	59 (60.8)	61 (47.3)	68 (59.6)	24 (51.1)
Injection drug user (IDU)	16 (11.5)	19 (15.1)	12 (12.4)	14 (10.9)	13 (11.4)	8 (17.0)
Gay or bisexual male IDU	20 (14.4)	23 (18.3)	16 (16.5)	29 (22.5)	21 (18.4)	11 (23.4)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	7 (5.0)	11 (8.7)	10 (10.3)	19 (14.7)	8 (7.0)	2 (4.3)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	0 (0.0)	1 (0.8)	0 (0.0)	6 (4.7)	4 (3.5)	2 (4.3)
Total	139 (100)	126 (100)	97 (100)	129 (100)	114 (100)	47 (100)

- (1) Data in recent years are incomplete due to delays in case reporting.
(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2019

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	657	349	13314
2003	650	332	13632
2004	691	340	13983
2005	565	348	14200
2006	538	326	14412
2007	535	324	14623
2008	520	263	14880
2009	472	251	15101
2010	470	244	15327
2011	426	236	15517
2012	466	236	15747
2013	399	259	15887
2014	322	240	15969
2015	281	256	15994
2016	235	236	15993
2017	228	254	15967
2018	199	210	15956
2019	81	80	15957

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2019

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	8	21
1982	99	32	88
1983	274	111	251
1984	557	273	535
1985	859	534	860
1986	1237	807	1290
1987	1636	878	2048
1988	1763	1038	2773
1989	2159	1279	3653
1990	2042	1364	4331
1991	2284	1512	5103
1992	2327	1639	5791
1993	2065	1603	6253
1994	1777	1600	6430
1995	1555	1485	6500
1996	1069	994	6575
1997	795	424	6946
1998	689	401	7234
1999	575	353	7456
2000	552	350	7658
2001	509	325	7842
2002	485	322	8005
2003	548	297	8256
2004	480	306	8430
2005	476	312	8594
2006	451	288	8757
2007	444	270	8931
2008	435	227	9139
2009	324	208	9255
2010	299	193	9361
2011	252	190	9423
2012	243	182	9484
2013	190	191	9483
2014	139	189	9433
2015	126	203	9356
2016	97	187	9266
2017	129	203	9192
2018	114	177	9129
2019	47	58	9118
Total	30131	21013	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Multiple races No. (%)	Total (1) No. (%)
Men						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	1 (0.0)	0 (0.0)	1 (0.2)	2 (0.0)
18 - 24	15 (0.2)	15 (1.0)	23 (0.8)	6 (0.6)	3 (0.7)	62 (0.4)
25 - 29	79 (0.9)	50 (3.5)	107 (3.8)	44 (4.6)	8 (1.8)	289 (2.0)
30 - 39	608 (6.8)	156 (10.9)	493 (17.4)	187 (19.6)	66 (14.6)	1510 (10.3)
40 - 49	1418 (15.8)	212 (14.8)	681 (24.0)	283 (29.6)	121 (26.7)	2717 (18.6)
50 - 59	3410 (38.1)	489 (34.1)	993 (35.0)	286 (29.9)	154 (34.0)	5334 (36.4)
60 +	3424 (38.2)	512 (35.7)	539 (19.0)	149 (15.6)	100 (22.1)	4726 (32.3)
Men Total	8954 (100)	1434 (100)	2837 (100)	955 (100)	453 (100)	14640 (100)
Women						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	1 (0.3)	1 (0.5)	0 (0.0)	0 (0.0)	2 (0.2)
18 - 24	2 (0.8)	6 (1.8)	5 (2.4)	0 (0.0)	2 (5.9)	15 (1.7)
25 - 29	5 (2.0)	7 (2.1)	10 (4.9)	1 (1.4)	2 (5.9)	25 (2.8)
30 - 39	25 (9.9)	31 (9.3)	37 (18.0)	11 (15.1)	4 (11.8)	108 (12.0)
40 - 49	48 (19.0)	56 (16.7)	41 (20.0)	22 (30.1)	9 (26.5)	177 (19.6)
50 - 59	103 (40.9)	119 (35.5)	55 (26.8)	25 (34.2)	12 (35.3)	314 (34.9)
60 +	69 (27.4)	115 (34.3)	56 (27.3)	14 (19.2)	5 (14.7)	260 (28.9)
Women Total	252 (100)	335 (100)	205 (100)	73 (100)	34 (100)	901 (100)
Trans Women	74 (100)	124 (100)	147 (100)	43 (100)	21 (100)	409 (100)

(1) Total includes persons with unknown race. Data on trans men are not released separately due to small numbers.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2019

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White		African American		Latino		Asian/ Pacific Islander/ Native American		Multiple races		Total (2)	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Men												
Gay or bisexual male	7317	(81.7)	867	(60.5)	2331	(82.2)	818	(85.7)	323	(71.3)	11658	(79.6)
Injection drug user (IDU)	175	(2.0)	185	(12.9)	72	(2.5)	21	(2.2)	16	(3.5)	469	(3.2)
Gay or bisexual male IDU	1376	(15.4)	276	(19.2)	336	(11.8)	78	(8.2)	103	(22.7)	2169	(14.8)
Transfusion/Hemophiliac	6	(0.1)	2	(0.1)	2	(0.1)	3	(0.3)	1	(0.3)	14	(0.2)
Heterosexual (3)	33	(0.4)	71	(5.0)	52	(1.8)	21	(2.2)	6	(1.3)	183	(1.3)
Perinatal	1	(0.0)	3	(0.2)	2	(0.1)	1	(0.1)	2	(0.4)	9	(0.1)
Other/Unknown	46	(0.5)	30	(2.1)	42	(1.5)	13	(1.4)	2	(0.4)	138	(0.9)
Men Total	8954	(100)	1434	(100)	2837	(100)	955	(100)	453	(100)	14640	(100)
Women												
Injection drug user (IDU)	146	(57.9)	164	(49.0)	62	(30.2)	13	(17.8)	23	(67.6)	408	(45.3)
Transfusion/Hemophilia	5	(2.0)	1	(0.3)	2	(1.0)	2	(2.7)	0	(0.0)	10	(1.1)
Heterosexual (3)	82	(32.5)	139	(41.5)	112	(54.6)	51	(69.9)	6	(17.6)	390	(43.3)
Perinatal	1	(0.4)	5	(1.5)	9	(4.4)	0	(0.0)	3	(8.8)	18	(2.0)
Other/Unknown	18	(7.1)	26	(7.8)	20	(9.8)	7	(9.6)	2	(5.9)	75	(8.3)
Women Total	252	(100)	335	(100)	205	(100)	73	(100)	34	(100)	901	(100)
Trans Women	74	(100)	124	(100)	147	(100)	43	(100)	21	(100)	409	(100)

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Total includes persons with unknown race. Data on trans men are not released separately due to small numbers.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.