

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through June 2020

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (628) 217-6336, FAX (415) 431-0353

Director of Health: Grant Colfax, MD
Director of Population Health Division: Tomas Aragon, MD, DrPH
Director of ARCHES: Wayne Enanoria, PhD, MPH
Director of HIV Surveillance Section: Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco^a (as of 06/30/2020)

Persons living with HIV infection (all disease stages)	15,898
Persons living with HIV infection ever classified as stage 3 (AIDS)	9,012
Diagnoses of HIV infection, 2018	205
Deaths of persons diagnosed with HIV infection, 2018	257
Deaths of persons diagnosed with HIV infection ever classified as stage 3 (AIDS), 2018	209

California^b (as of 12/31/2018)

Persons living with HIV infection (all disease stages)	136,566
Persons living with HIV infection ever classified as stage 3 (AIDS) ^c	66,887
Diagnoses of HIV infection, 2018	4,747
Deaths of persons diagnosed with HIV infection, 2018	1,872
Deaths of persons diagnosed with HIV infection ever classified as stage 3 (AIDS), 2018 ^d	1,395

United States^e (as of 12/31/2018)

Persons living with HIV infection (all disease stages)	1,042,270
Persons living with HIV infection ever classified as stage 3 (AIDS)	NA
Diagnoses of HIV infection, 2018	37,968
Deaths of persons diagnosed with HIV infection, 2018	15,821
Deaths of persons diagnosed with HIV infection ever classified as stage 3 (AIDS), 2018 ^d	12,373

^a Residents of San Francisco at time of HIV or AIDS diagnosis.


^b California HIV Surveillance Report 2018.
https://www.cdph.ca.gov/Programs/CID/DOA/Pages/OA_case_surveillance_reports.aspx. Released 2/27/2020.

^c Supplemental Tables to the California HIV Surveillance Report — 2018.
https://www.cdph.ca.gov/Programs/CID/DOA/Pages/OA_case_surveillance_reports.aspx. Released 4/13/2020.

^d CDC. Monitoring selected national HIV prevention and care objectives by using HIV surveillance data—United States and 6 dependent areas, 2018. HIV Surveillance Supplemental Report 2020;25(No. 2). (Table 7f)
<http://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>. Published May 2020.


^e CDC. HIV Surveillance Report, 2018 (Updated); vol. 31. (Table 1b, 10b, 14b).
<http://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>. Published May 2020.

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2020*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2020 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2009-2020

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	3 (0.6)	0 (0.0)	0 (0.0)	1 (0.2)	1 (0.3)
18 - 24	64 (13.5)	61 (12.9)	46 (10.6)	64 (13.7)	61 (15.1)	39 (12.1)
25 - 29	66 (13.9)	66 (14.0)	69 (16.0)	75 (16.0)	80 (19.8)	58 (18.0)
30 - 39	141 (29.7)	146 (30.9)	119 (27.5)	141 (30.1)	114 (28.2)	96 (29.8)
40 - 49	122 (25.7)	129 (27.3)	131 (30.3)	132 (28.2)	100 (24.8)	74 (23.0)
50 - 59	64 (13.5)	51 (10.8)	52 (12.0)	44 (9.4)	35 (8.7)	41 (12.7)
60 +	17 (3.6)	16 (3.4)	15 (3.5)	12 (2.6)	13 (3.2)	13 (4.0)
Total	475 (100)	472 (100)	432 (100)	468 (100)	404 (100)	322 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	3 (1.0)	0 (0.0)	3 (1.3)	1 (0.5)	0 (0.0)	0 (0.0)
18 - 24	38 (13.1)	34 (14.3)	25 (10.6)	31 (15.1)	16 (9.6)	6 (10.0)
25 - 29	72 (24.7)	53 (22.4)	42 (17.8)	39 (19.0)	23 (13.9)	9 (15.0)
30 - 39	87 (29.9)	77 (32.5)	80 (33.9)	65 (31.7)	65 (39.2)	19 (31.7)
40 - 49	60 (20.6)	37 (15.6)	49 (20.8)	42 (20.5)	28 (16.9)	15 (25.0)
50 - 59	18 (6.2)	24 (10.1)	26 (11.0)	17 (8.3)	19 (11.4)	8 (13.3)
60 +	13 (4.5)	12 (5.1)	11 (4.7)	10 (4.9)	15 (9.0)	3 (5.0)
Total	291 (100)	237 (100)	236 (100)	205 (100)	166 (100)	60 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2009-2020

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	15 (4.6)	15 (5.0)	12 (4.8)	11 (4.5)	10 (5.3)	5 (3.6)
25 - 29	23 (7.1)	26 (8.7)	21 (8.3)	25 (10.3)	17 (9.0)	13 (9.4)
30 - 39	75 (23.1)	74 (24.7)	62 (24.6)	60 (24.7)	43 (22.8)	41 (29.5)
40 - 49	120 (37.0)	99 (33.1)	99 (39.3)	83 (34.2)	60 (31.7)	40 (28.8)
50 - 59	69 (21.3)	62 (20.7)	49 (19.4)	44 (18.1)	38 (20.1)	27 (19.4)
60 +	21 (6.5)	23 (7.7)	9 (3.6)	20 (8.2)	21 (11.1)	13 (9.4)
Total	324 (100)	299 (100)	252 (100)	243 (100)	189 (100)	139 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	1 (0.8)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	4 (3.2)	2 (2.1)	1 (0.8)	3 (2.6)	2 (1.8)	0 (0.0)
25 - 29	9 (7.1)	8 (8.2)	15 (11.5)	12 (10.5)	9 (8.2)	4 (12.5)
30 - 39	35 (27.8)	27 (27.8)	31 (23.8)	32 (28.1)	33 (30.0)	11 (34.4)
40 - 49	36 (28.6)	23 (23.7)	41 (31.5)	21 (18.4)	25 (22.7)	3 (9.4)
50 - 59	28 (22.2)	19 (19.6)	15 (11.5)	26 (22.8)	23 (20.9)	6 (18.8)
60 +	14 (11.1)	18 (18.6)	26 (20.0)	20 (17.5)	18 (16.4)	8 (25.0)
Total	126 (100)	97 (100)	130 (100)	114 (100)	110 (100)	32 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2009-2020

Gender (2)	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Men	422 (88.8)	419 (88.8)	377 (87.3)	432 (92.3)	370 (91.6)	297 (92.2)
Women	31 (6.5)	36 (7.6)	42 (9.7)	24 (5.1)	22 (5.4)	13 (4.0)
Trans Women	22 (4.6)	17 (3.6)	13 (3.0)	12 (2.6)	12 (3.0)	12 (3.7)
Total	475 (100)	472 (100)	432 (100)	468 (100)	404 (100)	322 (100)

Gender (2)	Year of initial HIV diagnosis					
	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)
Men	255 (87.6)	202 (85.2)	203 (86.0)	180 (87.8)	137 (82.5)	49 (81.7)
Women	29 (10.0)	25 (10.5)	27 (11.4)	17 (8.3)	14 (8.4)	9 (15.0)
Trans Women	7 (2.4)	9 (3.8)	6 (2.5)	7 (3.4)	13 (7.8)	2 (3.3)
Total	291 (100)	237 (100)	236 (100)	205 (100)	166 (100)	60 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Data on trans men are not released separately due to small numbers.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2009-2020

Gender (2)	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Men	288 (88.9)	267 (89.3)	214 (84.9)	210 (86.4)	166 (87.8)	125 (89.9)
Women	24 (7.4)	16 (5.4)	24 (9.5)	24 (9.9)	17 (9.0)	7 (5.0)
Trans Women	12 (3.7)	16 (5.4)	14 (5.6)	9 (3.7)	6 (3.2)	7 (5.0)
Total	324 (100)	299 (100)	252 (100)	243 (100)	189 (100)	139 (100)

Gender (2)	Year of AIDS diagnosis					
	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)
Men	99 (78.6)	82 (84.5)	102 (78.5)	99 (86.8)	96 (87.3)	27 (84.4)
Women	19 (15.1)	11 (11.3)	22 (16.9)	12 (10.5)	7 (6.4)	4 (12.5)
Trans Women	8 (6.3)	3 (3.1)	6 (4.6)	3 (2.6)	7 (6.4)	1 (3.1)
Total	126 (100)	97 (100)	130 (100)	114 (100)	110 (100)	32 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Data on trans men are not released separately due to small numbers.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2009-2020

Race/ethnicity	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
White	227 (47.8)	225 (47.7)	221 (51.2)	230 (49.1)	181 (44.8)	135 (41.9)
Black/African American	61 (12.8)	65 (13.8)	63 (14.6)	51 (10.9)	51 (12.6)	32 (9.9)
Latino	117 (24.6)	109 (23.1)	89 (20.6)	113 (24.1)	106 (26.2)	95 (29.5)
Asian/Pacific Islander/ Native American	43 (9.1)	42 (8.9)	38 (8.8)	56 (12.0)	51 (12.6)	45 (14.0)
Multiple races	26 (5.5)	30 (6.4)	20 (4.6)	17 (3.6)	14 (3.5)	15 (4.7)
Total (2)	475 (100)	472 (100)	432 (100)	468 (100)	404 (100)	322 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)
White	119 (40.9)	90 (38.0)	83 (35.2)	60 (29.3)	58 (34.9)	14 (23.3)
Black/African American	43 (14.8)	34 (14.3)	39 (16.5)	42 (20.5)	28 (16.9)	14 (23.3)
Latino	77 (26.5)	70 (29.5)	70 (29.7)	79 (38.5)	54 (32.5)	22 (36.7)
Asian/Pacific Islander/ Native American	36 (12.4)	37 (15.6)	32 (13.6)	19 (9.3)	23 (13.9)	10 (16.7)
Multiple races	15 (5.2)	6 (2.5)	12 (5.1)	5 (2.4)	3 (1.8)	0 (0.0)
Total (2)	291 (100)	237 (100)	236 (100)	205 (100)	166 (100)	60 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2009-2020

Race/ethnicity	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
White	170 (52.5)	145 (48.5)	128 (50.8)	121 (49.8)	93 (49.2)	52 (37.4)
Black/African American	60 (18.5)	49 (16.4)	46 (18.3)	40 (16.5)	31 (16.4)	28 (20.1)
Latino	66 (20.4)	64 (21.4)	48 (19.0)	54 (22.2)	36 (19.0)	32 (23.0)
Asian/Pacific Islander/ Native American	15 (4.6)	23 (7.7)	14 (5.6)	14 (5.8)	18 (9.5)	17 (12.2)
Multiple races	13 (4.0)	18 (6.0)	16 (6.3)	14 (5.8)	11 (5.8)	10 (7.2)
Total (2)	324 (100)	299 (100)	252 (100)	243 (100)	189 (100)	139 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)
White	61 (48.4)	44 (45.4)	52 (40.0)	36 (31.6)	45 (40.9)	11 (34.4)
Black/African American	19 (15.1)	16 (16.5)	25 (19.2)	18 (15.8)	22 (20.0)	8 (25.0)
Latino	32 (25.4)	21 (21.6)	35 (26.9)	41 (36.0)	21 (19.1)	7 (21.9)
Asian/Pacific Islander/ Native American	5 (4.0)	11 (11.3)	13 (10.0)	10 (8.8)	15 (13.6)	5 (15.6)
Multiple races	9 (7.1)	5 (5.2)	5 (3.8)	9 (7.9)	7 (6.4)	1 (3.1)
Total (2)	126 (100)	97 (100)	130 (100)	114 (100)	110 (100)	32 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2009-2020

Transmission category (2)	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Gay or bisexual male	329 (69.3)	308 (65.3)	305 (70.6)	361 (77.1)	313 (77.5)	242 (75.2)
Injection drug user (IDU)	26 (5.5)	39 (8.3)	29 (6.7)	21 (4.5)	21 (5.2)	25 (7.8)
Gay or bisexual male IDU	81 (17.1)	69 (14.6)	61 (14.1)	51 (10.9)	48 (11.9)	42 (13.0)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	26 (5.5)	38 (8.1)	30 (6.9)	27 (5.8)	17 (4.2)	11 (3.4)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	13 (2.7)	18 (3.8)	7 (1.6)	8 (1.7)	5 (1.2)	2 (0.6)
Total	475 (100)	472 (100)	432 (100)	468 (100)	404 (100)	322 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)
Gay or bisexual male	220 (75.6)	166 (70.0)	145 (61.4)	130 (63.4)	111 (66.9)	40 (66.7)
Injection drug user (IDU)	17 (5.8)	21 (8.9)	25 (10.6)	27 (13.2)	12 (7.2)	3 (5.0)
Gay or bisexual male IDU	25 (8.6)	26 (11.0)	33 (14.0)	27 (13.2)	20 (12.0)	4 (6.7)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	21 (7.2)	17 (7.2)	16 (6.8)	11 (5.4)	9 (5.4)	2 (3.3)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	8 (2.7)	7 (3.0)	17 (7.2)	10 (4.9)	14 (8.4)	11 (18.3)
Total	291 (100)	237 (100)	236 (100)	205 (100)	166 (100)	60 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2009-2020

Transmission category (2)	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Gay or bisexual male	208 (64.2)	170 (56.9)	148 (58.7)	151 (62.1)	112 (59.3)	95 (68.3)
Injection drug user (IDU)	35 (10.8)	23 (7.7)	34 (13.5)	23 (9.5)	19 (10.1)	16 (11.5)
Gay or bisexual male IDU	56 (17.3)	83 (27.8)	49 (19.4)	43 (17.7)	45 (23.8)	21 (15.1)
Transfusion/Hemophilia	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	18 (5.6)	18 (6.0)	17 (6.7)	21 (8.6)	12 (6.3)	7 (5.0)
Perinatal	1 (0.3)	0 (0.0)	1 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	5 (1.5)	5 (1.7)	3 (1.2)	5 (2.1)	1 (0.5)	0 (0.0)
Total	324 (100)	299 (100)	252 (100)	243 (100)	189 (100)	139 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)
Gay or bisexual male	72 (57.1)	58 (59.8)	62 (47.7)	68 (59.6)	61 (55.5)	15 (46.9)
Injection drug user (IDU)	19 (15.1)	12 (12.4)	14 (10.8)	13 (11.4)	13 (11.8)	4 (12.5)
Gay or bisexual male IDU	23 (18.3)	16 (16.5)	29 (22.3)	21 (18.4)	27 (24.5)	9 (28.1)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	11 (8.7)	11 (11.3)	19 (14.6)	8 (7.0)	5 (4.5)	1 (3.1)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	1 (0.8)	0 (0.0)	6 (4.6)	4 (3.5)	4 (3.6)	3 (9.4)
Total	126 (100)	97 (100)	130 (100)	114 (100)	110 (100)	32 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2020

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	656	348	13299
2003	649	332	13616
2004	692	336	13972
2005	566	346	14192
2006	542	326	14408
2007	537	324	14621
2008	524	263	14882
2009	475	251	15106
2010	472	244	15334
2011	432	236	15530
2012	468	235	15763
2013	404	259	15908
2014	322	239	15991
2015	291	256	16026
2016	237	235	16028
2017	236	254	16010
2018	205	257	15958
2019	166	206	15918
2020	60	80	15898

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2020

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	8	21
1982	99	32	88
1983	274	111	251
1984	557	273	535
1985	859	534	860
1986	1237	807	1290
1987	1636	878	2048
1988	1762	1038	2772
1989	2159	1279	3652
1990	2043	1364	4331
1991	2284	1512	5103
1992	2326	1639	5790
1993	2064	1603	6251
1994	1776	1600	6427
1995	1551	1485	6493
1996	1066	993	6566
1997	795	424	6937
1998	686	401	7222
1999	574	353	7443
2000	552	349	7646
2001	507	325	7828
2002	481	321	7988
2003	544	296	8236
2004	477	300	8413
2005	476	310	8579
2006	451	288	8742
2007	442	270	8914
2008	435	227	9122
2009	324	208	9238
2010	299	193	9344
2011	252	190	9406
2012	243	181	9468
2013	189	191	9466
2014	139	188	9417
2015	126	203	9340
2016	97	186	9251
2017	130	203	9178
2018	114	209	9083
2019	110	149	9044
2020	32	64	9012
Total	30197	21185	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	Black/African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Multiple races No. (%)	Total (1) No. (%)
Men						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.2)	1 (0.0)
18 - 24	8 (0.1)	13 (0.9)	22 (0.8)	6 (0.6)	1 (0.2)	50 (0.3)
25 - 29	61 (0.7)	46 (3.2)	99 (3.5)	41 (4.3)	13 (2.8)	260 (1.8)
30 - 39	558 (6.3)	161 (11.3)	491 (17.1)	186 (19.3)	57 (12.3)	1454 (10.0)
40 - 49	1264 (14.3)	198 (13.8)	639 (22.3)	268 (27.9)	112 (24.2)	2483 (17.0)
50 - 59	3277 (37.1)	475 (33.2)	1013 (35.3)	291 (30.2)	167 (36.1)	5224 (35.9)
60 +	3670 (41.5)	538 (37.6)	605 (21.1)	170 (17.7)	111 (24.0)	5097 (35.0)
Men Total	8838 (100)	1431 (100)	2869 (100)	962 (100)	462 (100)	14569 (100)
Women						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	1 (0.3)	1 (0.5)	0 (0.0)	0 (0.0)	2 (0.2)
18 - 24	1 (0.4)	5 (1.5)	1 (0.5)	0 (0.0)	2 (6.1)	9 (1.0)
25 - 29	4 (1.6)	7 (2.1)	12 (5.8)	1 (1.3)	2 (6.1)	26 (2.9)
30 - 39	25 (10.0)	34 (10.2)	34 (16.5)	8 (10.7)	4 (12.1)	105 (11.7)
40 - 49	46 (18.4)	53 (15.9)	44 (21.4)	20 (26.7)	9 (27.3)	173 (19.2)
50 - 59	102 (40.8)	107 (32.0)	59 (28.6)	26 (34.7)	12 (36.4)	306 (34.0)
60 +	72 (28.8)	127 (38.0)	55 (26.7)	20 (26.7)	4 (12.1)	279 (31.0)
Women Total	250 (100)	334 (100)	206 (100)	75 (100)	33 (100)	900 (100)
Trans Women	78 (100)	129 (100)	151 (100)	42 (100)	21 (100)	421 (100)

(1) Total includes persons with unknown race. Data on trans men are not released separately due to small numbers.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2020

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White No. (%)	Black/African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Multiple races No. (%)	Total (2) No. (%)
Men						
Gay or bisexual male	7234 (81.9)	876 (61.2)	2366 (82.5)	821 (85.3)	332 (71.9)	11631 (79.8)
Injection drug user (IDU)	169 (1.9)	176 (12.3)	70 (2.4)	21 (2.2)	16 (3.5)	452 (3.1)
Gay or bisexual male IDU	1350 (15.3)	269 (18.8)	334 (11.6)	82 (8.5)	104 (22.5)	2139 (14.7)
Transfusion/Hemophiliac	5 (0.1)	2 (0.1)	2 (0.1)	3 (0.3)	0 (0.3)	12 (0.0)
Heterosexual (3)	32 (0.4)	71 (5.0)	51 (1.8)	20 (2.1)	6 (1.3)	180 (1.2)
Perinatal	1 (0.0)	3 (0.2)	2 (0.1)	1 (0.1)	2 (0.4)	9 (0.1)
Other/Unknown	47 (0.5)	34 (2.4)	44 (1.5)	14 (1.5)	2 (0.4)	146 (1.0)
Men Total	8838 (100)	1431 (100)	2869 (100)	962 (100)	462 (100)	14569 (100)
Women						
Injection drug user (IDU)	141 (56.4)	161 (48.2)	60 (29.1)	13 (17.3)	22 (66.7)	397 (44.1)
Transfusion/Hemophilia	5 (2.0)	1 (0.3)	2 (1.0)	2 (2.7)	0 (0.0)	10 (1.1)
Heterosexual (3)	83 (33.2)	138 (41.3)	113 (54.9)	52 (69.3)	6 (18.2)	392 (43.6)
Perinatal	1 (0.4)	6 (1.8)	9 (4.4)	0 (0.0)	2 (6.1)	18 (2.0)
Other/Unknown	20 (8.0)	28 (8.4)	22 (10.7)	8 (10.7)	3 (9.1)	83 (9.2)
Women Total	250 (100)	334 (100)	206 (100)	75 (100)	33 (100)	900 (100)
Trans Women	78 (100)	129 (100)	151 (100)	42 (100)	21 (100)	421 (100)

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Total includes persons with unknown race. Data on trans men are not released separately due to small numbers.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.